

AQA Physics (Separate Science) Unit 3: Particle Model of Matter

Required Practical

Measuring the density of a regularly shaped object:

- Measure the mass using a balance.
- Measure the length, width and height using a ruler.
- Calculate the volume.
- Use the density ($\rho = m/V$) equation to calculate density.

Measuring the density of an irregularly-shaped object:

- Measure the mass using a balance.
- Fill a eureka can with water.
- Place the object in the water - the water displaced by the object will transfer into a measuring cylinder.
- Measure the volume of the water. This equals the volume of the object.
- Use the density ($\rho = m/V$) equation to calculate density.



Density

Density is a measure of how much mass there is in a given space.

$$\text{Density (kg/m}^3\text{)} = \text{mass (kg)} \div \text{volume (m}^3\text{)}$$

A more dense material will have more particles in the same volume when compared to a less dense material.

Example

The density of an object is 8050kg/m^3 and it has a volume of 3.4m^3 .

What is its mass in kg?


$$8050 = \text{mass} \div 3.4$$

$$8050 \times 3.4 = \text{mass}$$


$$27\,370\text{kg}$$

States of Matter


Solids have strong forces of attraction between the particles. The particles are held together very closely in a fixed, regular arrangement. The particles do not have much energy and can only vibrate.



Liquids have weaker forces of attraction between the particles. The particles are close together, but can move past each other. They form irregular arrangements. They have more energy than particles in a solid.




Gases have almost no forces of attraction between the particles. The particles have the most energy and are free to move in random directions.




Motion in Gas Particles

Gas particles move about randomly, at high speed. They intercept other gas particles and anything else that is in the way. When this occurs, a pressure is exerted. If the gas is within a sealed container, pressure occurs when the gas particles hit the walls of the container. They produce a force at right angles to the wall of the container.




If the temperature of the gas increases, then the pressure will also increase. The hotter the temperature, the more kinetic energy the gas particles have. They move faster, colliding with the sides of the container more often.




Internal Energy

Particles within a system have kinetic energy when they vibrate or move around. The particles also have a potential energy store. The total internal energy of a system is the kinetic and potential energy stores.




Low Temperature



High Temperature

If the system is heated, the particles will gain more kinetic energy, so increasing the internal energy.

Changing State




If a system gains more energy, it can lead to a change in temperature or change in state. If the system is heated enough, then there will be enough energy to break bonds.


When something changes state, there is no chemical change, only physical. No new substance is formed. The substance will change back to its original form. The number of particles does not change and mass is conserved.

Specific Latent Heat

Energy is being put in during melting and boiling. This increases the amount of internal energy. The energy is being used to break the bonds, so the temperature does not increase. This is shown by the parts of the graph that are flat.



When a substance is condensing or freezing, the energy put in is used to form the bonds. This releases energy. The internal energy decreases, but the temperature does not go down.



The energy needed to change the state of a substance is called the latent heat.

Specific latent heat is the amount of energy needed to change 1kg of a substance from one state to another without changing the temperature. Specific latent heat will be different for different materials.

- solid → liquid - specific latent heat of **fusion**
- liquid → gas - specific latent heat of **vaporisation**

Specific Latent Heat Equation

The amount of energy needed/released when a substance of mass changes state.

$$\text{energy (E)} = \text{mass (m)} \times \text{specific latent heat (L)}$$

$$E = mL$$



Pressure in Gases

Heating

Heating up the gas particles provides them with more energy to move more quickly. This means they are likely to collide more frequently with other particles. This, in turn, increases the pressure.

Volume

If the volume of the container is increased, the number of collisions will decrease. This causes an overall decrease in pressure.

The equation for a fixed mass and a constant temperature is as follows (you will be given this in the exam):

$$P \times V = \text{constant}$$

P = pressure (Pa)

V = volume (m³)

If the volume increases, the pressure decreases.

If the volume decreases, the pressure increases.

If the pressure of a gas changes, the volume of the gas can also change. A helium-filled balloon, once released, will rise into the atmosphere; the pressure outside of the balloon will decrease. The volume of the balloon will increase (due to less pressure outside the balloon), meaning the pressure inside the balloon will decrease.

Work Done on a Gas (Higher Tier Only)

Work done on a gas causes it to gain internal energy and so will increase the temperature. Pumping up the tyre of a bicycle involves doing work and this will increase the temperature of the gas inside the bicycle tyre.

